

## PDF hosted at the Radboud Repository of the Radboud University Nijmegen

The following full text is a publisher's version.

For additional information about this publication click this link.

<http://hdl.handle.net/2066/207986>

Please be advised that this information was generated on 2019-11-08 and may be subject to change.

# Endovascular Treatment With or Without Prior Intravenous Alteplase for Acute Ischemic Stroke

Vicky Chalos, MD;\* Natalie E. LeCouffe, MD;\* Maarten Uyttenboogaart, MD, PhD; Hester F. Lingsma, PhD; Maxim J. H. L. Mulder, MD, PhD; Esmee Venema, MD; Kilian M. Treurniet, MD; Omid Eshghi, MD; H. Bart van der Worp, MD, PhD; Aad van der Lugt, MD, PhD; Yvo B. W. E. M. Roos, MD, PhD; Charles B. L. M. Majoie, MD, PhD; Diederik W. J. Dippel, MD, PhD; Bob Roozenbeek, MD, PhD;\*\* Jonathan M. Coutinho, MD, PhD\*\*\*; on behalf of the MR CLEAN Registry Investigators\*\*\*

**Background**—It is unclear whether intravenous thrombolysis (IVT) with alteplase before endovascular treatment (EVT) is beneficial for patients with acute ischemic stroke caused by a large vessel occlusion. We compared clinical and procedural outcomes, safety, and workflow between patients treated with both IVT and EVT and those treated with EVT alone in routine clinical practice.

**Methods and Results**—Using multivariable regression, we evaluated the association of IVT+EVT with 90-day functional outcome (modified Rankin Scale), mortality, reperfusion, first-pass effect, and symptomatic intracranial hemorrhage in the MR CLEAN (Multicenter Randomised Controlled Trial of Endovascular Treatment for Acute Ischemic Stroke in The Netherlands) Registry. Of 1485 patients, 1161 (78%) were treated with IVT+EVT, and 324 (22%) with EVT alone. Patients treated with IVT+EVT had atrial fibrillation less often (16% versus 44%) and had better pre-stroke modified Rankin Scale scores (pre-stroke modified Rankin Scale 0: 73% versus 52%) than those treated with EVT alone. Procedure time was shorter in the IVT+EVT group (median 62 versus 68 minutes). Nontransferred IVT+EVT patients had longer door-to-groin-puncture times (median 105 versus 94 minutes). IVT+EVT was associated with better functional outcome (adjusted common odds ratio 1.47; 95% CI: 1.10–1.96) and lower mortality (adjusted odds ratio 0.58; 95% CI: 0.40–0.82). Successful reperfusion, first-pass effect, and symptomatic intracranial hemorrhage did not differ between groups.

**Conclusions**—In this observational study, patients treated with IVT+EVT had better clinical outcomes than patients who received EVT alone. This finding may demonstrate a true benefit of IVT before EVT, but its interpretation is hampered by the possibility of residual confounding and selection bias. Randomized trials are required to properly assess the effect of IVT before EVT. (*J Am Heart Assoc.* 2019;8:e011592. DOI: 10.1161/JAHA.118.011592.)

**Key Words:** endovascular treatment • large vessel occlusion • stroke • thrombectomy • thrombolysis

Endovascular treatment (EVT) has become standard of care for patients with acute ischemic stroke caused by an intracranial large vessel occlusion of the anterior circulation.<sup>1</sup> Patients included in the EVT trials received intravenous thrombolysis (IVT) with alteplase as standard care, unless they had a contraindication for IVT. Hence, all major guidelines recommend

IVT in eligible patients before EVT.<sup>2</sup> In a recent meta-analysis of randomized trials, the effect of EVT was not influenced by IVT, raising the question of whether IVT is beneficial to patients with a large vessel occlusion.<sup>1</sup> Theoretical advantages of IVT before EVT include early reperfusion, faster procedural times, lysis of distal emboli, and improved microvascular reperfusion.<sup>3</sup> Potential

From the Department of Neurology (V.C., M.J.H.L.M., D.W.J.D., B.R.), Public Health, Center for Medical Decision Making (V.C., H.F.L., E.V.), and Radiology and Nuclear Medicine (V.C., A.v.d.L., B.R.), Erasmus MC University Medical Center Rotterdam, Rotterdam, The Netherlands; Department of Neurology (N.E.L., Y.B.W.E.M.R., J.M.C.) and Radiology and Nuclear Medicine (N.E.L., K.M.T., C.B.L.M.M.), Amsterdam UMC, University of Amsterdam, The Netherlands; Department of Neurology (M.U.) and Radiology (M.U., O.E.), University Medical Center Groningen, Groningen, The Netherlands; Department of Neurology and Neurosurgery, Brain Center Rudolf Magnus, University Medical Center Utrecht, Utrecht, The Netherlands (H.B.v.d.W.).

\*Dr Chalos and Dr LeCouffe contributed equally to this work as co-first authors.

\*\*Dr Roozenbeek and Dr Coutinho contributed equally to this work as co-last authors.

\*\*\*A complete list of the MR CLEAN Registry Investigators can be found in the Appendix at the end of the article.

**Correspondence to:** Jonathan M. Coutinho, MD, PhD, Department of Neurology, Amsterdam UMC, University of Amsterdam, Meibergdreef 9, P.O. Box 22660 1100 DD Amsterdam, The Netherlands. E-mail: j.coutinho@amc.uva.nl

Received December 5, 2018; accepted March 20, 2019.

© 2019 The Authors. Published on behalf of the American Heart Association, Inc., by Wiley. This is an open access article under the terms of the Creative Commons Attribution-NonCommercial-NoDerivs License, which permits use and distribution in any medium, provided the original work is properly cited, the use is non-commercial and no modifications or adaptations are made.

## Clinical Perspective

### What Is New?

- The question of whether intravenous thrombolysis (IVT) should still be administered before endovascular treatment (EVT) in ischemic stroke patients was investigated in a large, nationwide registry of patients who underwent EVT, reflecting daily clinical practice.
- IVT administration appeared to delay the time until groin puncture in nontransferred patients.
- Various aspects of this debate, including clinical and safety outcomes, workflow implications, first-pass effect (successful reperfusion on first pass without rescue medication), and reperfusion of the occluded vessel before EVT were explored in 1 comprehensive study.

### What Are the Clinical Implications?

- In addition to having better clinical outcomes, patients treated with IVT+EVT more often had reperfusion of the occluded vessel before EVT than patients treated with EVT alone.
- IVT remains the only available reperfusion therapy in about 6% of all patients, in whom the clot was not accessible because of technical reasons; however, successful reperfusion and first-pass effect did not differ between groups, implying that IVT does not facilitate the thrombectomy procedure.
- Until results of ongoing randomized trials are published, current guidelines should remain unchanged.

disadvantages include delayed initiation of EVT, thrombus fragmentation, major bleeding, potential neurotoxicity, and disruption of the blood–brain barrier.<sup>4–6</sup>

Several studies examined the efficacy and safety of IVT before EVT, but the sample sizes of most studies were relatively small, and their results were inconclusive.<sup>7–10</sup>

The aim of our study was to compare clinical and procedural outcomes, safety, and workflow between acute ischemic stroke patients with a large vessel occlusion treated with both IVT and EVT to those treated with EVT alone using data of the MR CLEAN (Multicenter Randomised Controlled Trial of Endovascular Treatment for Acute Ischemic Stroke in the Netherlands) Registry.

## Methods

### Data, Materials, and Code Disclosure Statement

Data will not be made available to other researchers for purposes of reproducing the results or replicating the procedure, because no patient approval has been obtained for sharing coded data. However, syntax and output files of statistical analyses may be made available upon request.

## Study Design

Details of the MR CLEAN Registry have been reported previously.<sup>11</sup> Briefly, the MR CLEAN Registry is an ongoing, nationwide, multicenter, prospective, observational phase IV study for centers that provide EVT in The Netherlands. Data are collected from consecutive patients who underwent EVT in 18 hospitals. All imaging of patients in the MR CLEAN Registry is adjudicated by an imaging core laboratory, whose members are blinded to clinical findings, except for symptom side. Safety parameters are scored by the complication committee, whose members are blinded to treatment center. A central medical ethics committee evaluated the study protocol of the MR CLEAN Registry and granted permission to carry out the study as a registry.

## Study Population and Treatment

We included adult patients who were treated in a MR CLEAN trial center between March 2014 and June 2016, with a large vessel occlusion of the anterior circulation confirmed on computed tomography angiography or magnetic resonance angiography (intracranial carotid artery [ICA/ICA-T], middle cerebral artery [M1/M2], anterior cerebral artery [A1/A2]), and who underwent groin puncture within 6.5 hours after symptom onset. We excluded patients for whom it was unknown whether they received IVT.

IVT (0.9 mg/kg alteplase over 1 hour with 10% initial bolus) was administered at the first hospital of arrival, according to national guidelines. EVT consisted of mechanical thrombectomy with a stent retriever and/or thrombus aspiration, with or without local delivery of a thrombolytic agent.

## Outcome Measures

The primary outcome was the modified Rankin Scale (mRS) score at 90 days. Secondary outcomes were an mRS score of 0 to 2 (functional independence) at 90 days; change in score on the National Institutes of Health Stroke Scale (NIHSS) from baseline to 24 to 48 hours (delta NIHSS); door-(intervention center)-to-groin-puncture time; procedure time; onset-to-last-contrast-bolus time; first-pass effect (single pass/use of the device as first line of EVT, complete reperfusion of the large vessel occlusion and its downstream territory [eTICI 3] and no use of rescue therapy after use of the device)<sup>12</sup>; reperfusion before start of EVT (defined as a score  $\geq 2B$  on the extended Thrombolysis in Cerebral Ischemia scale [eTICI] on first digital subtraction angiography); and successful reperfusion post-EVT (defined as an eTICI score  $\geq 2B$  or  $\geq 2C$ ).<sup>13</sup>

Procedure time was defined as the moment of puncture of the femoral artery to successful reperfusion (eTICI  $\geq 2B$ ) or last contrast bolus (when successful reperfusion was not achieved or no target occlusion was seen during the intervention).

Onset-to-last-contrast-bolus time was defined as the duration from symptom onset or time of last seen well to successful reperfusion or last contrast bolus. Safety outcomes were mortality at 90 days, severe extracranial hemorrhage (ie, requiring surgery or blood transfusion), and symptomatic intracranial hemorrhage (sICH) according to the Heidelberg Bleeding Classification.<sup>14</sup>

## Statistical Analysis

We compared patients treated with IVT+EVT with patients treated with EVT alone. For intergroup comparison, we used a  $\chi^2$  test, Fisher's exact test, Student *t* test, or Mann–Whitney *U* test. The mRS scores of patients treated with IVT+EVT were compared with those of patients treated with EVT alone by means of ordinal logistic regression. Binary outcomes were analyzed with logistic regression analysis and continuous outcomes with linear regression analysis. We made adjustments based on theoretical identification, known association with outcome, and empirical identification (ie, baseline imbalances). For all analyses, we made adjustments in a multivariable model for age, baseline NIHSS, history of diabetes mellitus, pre-stroke mRS, prior use of anticoagulant medication, onset-to-first-noncontrast-computed-tomography time, center (in case of sufficient  $\geq 1$  outcome events), and additional baseline imbalances ( $P < 0.05$ ) in the patients' medical histories. For clinical outcome measures, we made additional adjustments for baseline mean arterial pressure, occlusion location, collateral score,<sup>15</sup> and transfer from a primary stroke center. For door-to-groin-puncture time, procedure time, onset-to-last-contrast-bolus time, first-pass effect, reperfusion before start of EVT, and successful reperfusion, we additionally adjusted for occlusion location and transfer from a primary stroke center. For sICH and severe extracranial hemorrhage, we additionally adjusted for baseline mean arterial pressure, prior use of antiplatelet agents, and Alberta Stroke Program Early CT Score. We tested for collinearity between all variables in all analyses by measuring the variance inflation factor.

In a supplementary analysis, we compared the distribution of occlusion locations in patients with reperfusion before start of EVT to those without.

Missing data were imputed using multiple imputation based on relevant covariates and outcome. Adjusted (common) odds ratios [a(c)ORs] are reported with 95% CIs and all *P* values are 2-sided. Statistical analyses were performed with Stata software, version 14.1 (StatCorp, TX), and IBM SPSS Statistics for Windows, Version 24.0.

## Sensitivity Analyses

We performed 2 sensitivity analyses. First, we conducted a 1:1 propensity score matching analysis to evaluate the

association between IVT and functional outcome. Propensity scores representing the probability of receiving IVT were calculated for each patient in each multiple imputed data set, using a logistic regression model, based on the covariates used for the adjustments in our primary analysis. Patients from the EVT alone group were matched to patients in the IVT+EVT group in a 1:1 nearest-neighbor matching of the logit of the propensity score, with a caliper width of 0.20. Matching was performed without replacement and unpaired patients were excluded. We used an ordinal logistic regression analysis to compare functional outcomes of patients treated with IVT+EVT and EVT alone.

To explore residual confounding, we performed a second sensitivity analysis, in which we stratified for “history of atrial fibrillation.” In this analysis, we used the outcome variables mRS at 90 days, functional independence at 90 days, mortality at 90 days, sICH, and eTICI  $\geq 2$ B.

## Results


During the study period, 1628 patients were recorded in the MR CLEAN Registry, of whom 140 did not meet the inclusion criteria. We further excluded 3 patients because it was unknown whether they had been treated with IVT. Of the 1485 included patients, 1161 (78%) were treated with IVT+EVT and 324 (22%) with EVT alone (Figure 1).

The most common reasons for withholding IVT were coagulation abnormalities and/or antithrombotic treatment (50%), recent surgery (15%), and presentation exceeding 4.5 hours after symptom onset or last seen well (14%) (Table 1).

Patients in the IVT+EVT group were younger (median 70 years versus 72 years,  $P = 0.03$ ), had less severe deficits (median NIHSS 16 versus 17,  $P < 0.01$ ), less often had atrial fibrillation (16% versus 44%,  $P < 0.01$ ) and previous ischemic stroke (14% versus 26%,  $P < 0.01$ ) than patients treated with EVT alone (Table 2). Patients treated with IVT+EVT also had a better pre-stroke mRS (mRS 0: 73% versus 52%,  $P < 0.01$ ).

In total, 656 (57%) patients with IVT+EVT were transferred from a primary stroke center, compared with 151 (47%) patients treated with EVT alone ( $P < 0.01$ ) (Table 3). The median door-to-needle time in patients treated with IVT+EVT was 25 minutes. Onset-to-noncontrast-computed-tomography time (median 67 versus 83 minutes,  $P < 0.01$ ) and onset-to-groin-puncture time (median 206 versus 215 minutes,  $P = 0.04$ ) were shorter in the IVT+EVT group.

The scores on the mRS at 90 days were more favorable in patients in the IVT+EVT group than in the EVT-alone group, and this difference persisted after adjustment for potential confounders (acOR 1.47; 95% CI: 1.10–1.96, Table 4, Figure 2). Functional independence at 90 days was achieved in 41% of patients with IVT+EVT, compared with 29% of patients with EVT



**Figure 1.** Flowchart of patient selection. EVT indicates endovascular treatment; IVT, intravenous thrombolysis; MR CLEAN, a Multicenter Randomized Controlled trial of Endovascular Treatment for Acute Ischemic Stroke in The Netherlands.

alone (aOR 1.32; 95% CI: 0.85–1.87). Delta NIHSS was larger (adjusted  $\beta$   $-1.5$ ; 95% CI:  $-2.6$  to  $-0.3$ ) and mortality was lower in the IVT+EVT group (aOR 0.58; 95% CI: 0.40–0.82).

Among nontransferred patients, median door-to-groin-puncture time was longer in the IVT+EVT group (105 versus 94 minutes; adjusted  $\beta$  9.5; 95% CI: 0.5–18.5), while among

transferred patients this was 47 minutes in both groups. We found no association between prior IVT and door-to-groin-puncture time for transferred patients only (adjusted  $\beta$  0.5; 95% CI:  $-7.3$  to  $8.2$ ), nor between prior IVT and onset-to-last-contrast-bolus time (adjusted  $\beta$   $-3.5$ ; 95% CI:  $-12.8$  to  $5.9$ ). There also was no difference in the proportion of patients with first-pass effect (17% versus 16%; aOR 1.22; 95% CI: 0.79–1.90). We did find a faster median procedure time (62 versus 68 minutes, adjusted  $\beta$   $-6.2$ ; 95% CI:  $-11.0$  to  $-1.3$ ) in the IVT+EVT group. Reperfusion before start of EVT occurred more often in the IVT+EVT group than in the EVT-alone group (8.4% versus 2.8%, aOR 3.14; 95% CI: 1.47–6.73) (Table 4).

In the supplementary analysis in 85 of the 97 IVT+EVT patients who had reperfusion before EVT and in whom the occlusion location was known, 61 (72%) had a distal M1, M2, or M3 occlusion (Table 5).

The risk of severe extracranial hemorrhage (2.4% versus 1.5%, aOR 1.96; 95% CI: 0.66–5.81) and sICH (5.9% versus 5.3%, aOR 1.20; 95% CI: 0.64–2.25) did not differ between groups (Table 4).

### Sensitivity Analyses

After applying 1:1 matching in each multiple imputed data set, sample sizes of both the IVT+EVT and EVT alone group ranged between 299 and 305. Baseline characteristics were similar in the 2 matched groups, with significant differences remaining in atrial fibrillation, prior use of direct oral anticoagulants, and prior use of vitamin K antagonists (Table 6). Prior IVT was still associated with better functional outcome at 90 days (cOR 1.42; 95% CI: 1.03–1.96).

**Table 1.** Reported Reasons for Withholding IVT

Reason for Withholding IVT*	Total (n=324), n (%)
Coagulation abnormalities and/or antithrombotic treatment	163 (50)
Recent surgery	48 (15)
Time from symptom onset or last seen well exceeds 4.5 h	46 (14)
Recent ischemic stroke	29 (9.0)
Recent traumatic injury or current hemorrhage	20 (6.2)
Recent gastrointestinal or urogenital hemorrhage	12 (3.7)
Other, such as allergy for IVT, cerebellar metastasis, endocarditis, pregnancy	10 (3.1)
Recent ICH	8 (2.5)
SBP $\geq 185$ mm Hg and/or DBP $\geq 110$ mm Hg	7 (2.2)
Unknown	5 (1.5)

DBP indicates diastolic blood pressure; ICH, intracranial hemorrhage; IVT, intravenous thrombolysis; SBP, systolic blood pressure.

\*More than 1 reason may have been reported per patient.



**Table 2.** Baseline Characteristics

	IVT+EVT (n=1161)	EVT (n=324)	P Value
Age (y), median (IQR)	70 (59–79)	72 (63–80)	0.03
Men, n (%)	621 (54)	171 (53)	0.82
NIHSS, median (IQR)*	16 (11–20)	17 (13–20)	<0.01
Systolic BP, mean mm Hg (SD) <sup>†</sup>	150 (24)	149 (26)	0.85
Diastolic BP, mean mm Hg (SD) <sup>‡</sup>	82 (15)	82 (17)	0.77
Medical history			
Atrial fibrillation, n (%)	186/1144 (16)	141/320 (44)	<0.01
Diabetes mellitus, n (%)	197/1155 (17)	56/321 (17)	0.87
Hypertension, n (%)	562/1145 (49)	180/321 (56)	0.03
Ischemic stroke, n (%)	164/1154 (14)	83/322 (26)	<0.01
Myocardial infarction, n (%)	163/1142 (14)	64/314 (20)	<0.01
Peripheral artery disease, n (%)	99/1139 (8.7)	36/318 (11)	0.15
Pre-stroke mRS, n (%)			<0.01
0	826/1138 (73)	165/320 (52)	
1	132/1138 (12)	57/320 (18)	
2	70/1138 (6.2)	38/320 (12)	
≥3	110/1138 (9.7)	60/320 (19)	
Medication			
Direct oral anticoagulants, n (%)	10/1141 (0.9)	27/318 (8.5)	<0.01
Vitamin K antagonists, n (%)	70/1150 (6.1)	120/324 (37)	<0.01
Antiplatelets, n (%)	391/1146 (34)	100/320 (31)	0.34
Imaging			
Occlusion location on CTA, n (%)			0.32
ICA	70/1101 (6.4)	12/307 (3.9)	
ICA-T	241/1101 (22)	71/307 (23)	
M1	637/1101 (58)	186/307 (61)	
M2	142/1101 (13)	33/307 (11)	
Other <sup>  </sup>	11/1101 (1.0)	5/307 (1.6)	
ASPECTS, median (IQR) <sup>§</sup>	9 (7–10)	9 (7–10)	0.76
Collateral score, n (%)			0.74
Grade 0	72/1105 (6.7)	24/303 (7.9)	
Grade 1	365/1105 (34)	95/303 (31)	
Grade 2	41/1105 (39)	117/303 (39)	
Grade 3	221/1105 (21)	67/303 (22)	

ASPECTS indicates Alberta Stroke Program Early CT Score; BP, blood pressure; CTA, computed tomography angiography; EVT, endovascular treatment; ICA, internal carotid artery; ICA-T, terminal internal carotid artery; IQR, interquartile range; IVT, intravenous thrombolysis; M1, middle cerebral artery segment 1; M2, middle cerebral artery segment 2; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale.

Missing: \*30; <sup>†</sup>42; <sup>‡</sup>47; <sup>§</sup>64.

<sup>||</sup>Other: occlusions in segment 1 or 2 of the anterior cerebral artery (A1: n=3; A2: n=3) or segment 3 of the middle cerebral artery (M3, n=9), or no occlusion visible (n=12) on CTA after adjudication by the imaging core laboratory.

We next stratified the analysis for atrial fibrillation. Among 327 patients with atrial fibrillation, 186 underwent IVT+EVT and 141 EVT alone. Of the 1137 patients without atrial fibrillation, 958 underwent IVT+EVT and 179 EVT alone. A

comparison of baseline characteristics within the strata is shown in Table 7. The association between prior IVT and better functional outcome at 90 days remained statistically significant among patients without atrial fibrillation (acOR

**Table 3.** Workflow\* and Treatment Characteristics

	IVT+EVT (n=1161)	EVT (n=324)	P Value
Transferred from primary stroke center, n (%)	656 (57)	151 (47)	<0.01
Onset-to-first-NCCT time <sup>†</sup>	67 (51–103)	83 (52–142)	<0.01
Door-to-needle time <sup>‡</sup>	25 (19–33)	NA	
Onset-to-groin-puncture time <sup>  </sup>	206 (160–260)	215 (158–294)	0.04
Door-to-groin-puncture time <sup>  §</sup>	128 (97–165)	115 (85–165)	0.08
Performed procedure, n (%)			
Catheterization—no access to target occlusion	64 (5.5)	16 (5.0)	<0.01
DSA—no target occlusion present	108 (9.3)	11 (3.4)	
Thrombectomy—thrombus retrieval attempted	983 (85)	294 (91)	
Other—procedure ended before attempt	6 (0.5)	3 (0.9)	

DSA indicates digital subtraction angiography; EVT, endovascular treatment; IQR, interquartile range; IVT, intravenous thrombolysis; NA, not applicable; NCCT, noncontrast computed tomography.

\*All times are in minutes—median (IQR).

Missing: <sup>†</sup>495; <sup>‡</sup>319; <sup>§</sup>447.

<sup>||</sup>Includes both transferred and nontransferred patients; door-time is door of first hospital.

1.72; 95% CI: 1.23–2.42, Table 8). However, among patients with atrial fibrillation, there was no association between prior IVT and functional outcome (acOR 1.06; 95% CI: 0.63–1.81). We also found a higher percentage of sICH in patients with atrial fibrillation who were treated with IVT+EVT, but this difference was not statistically significant (5.9% versus 2.8%, aOR 2.18; 95% CI: 0.60–7.91).

## Discussion

In our study—in which we compared clinical and procedural outcomes, safety, and workflow in patients with acute ischemic stroke and an intracranial large vessel occlusion of the anterior circulation treated with IVT+EVT to those treated with EVT alone—we found that the combination of EVT with IVT was associated with a better clinical outcome than treatment with EVT alone.

A number of observational studies have previously addressed the additional benefit of IVT before EVT. These studies mostly had small sample sizes (range: 66–500 patients, with the exception of 1 study of 1166 patients) and had varying results.<sup>7–9,16–18</sup> A post hoc, pooled analysis of the SWIFT (Solitaire With the Intention for Thrombectomy) and STAR (Solitaire Flow Restoration Thrombectomy for Acute Revascularization) studies showed no statistically significant benefit of IVT followed by EVT over EVT alone.<sup>9</sup> However, contrary to our study, the effect of IVT was awaited before initiating EVT in the majority of cases, possibly decreasing the chances of good functional outcome. Two other studies performed a propensity score matching analysis comparing patients who received IVT before EVT with IVT-eligible patients who underwent EVT alone, and also found no difference in

functional independence between the 2 groups.<sup>8,10</sup> In only 2 studies with data of 66 and 131 patients, a score on the mRS of 0 to 2 at 90 days was more common in the IVT+EVT group, which is in line with our results.<sup>16,17</sup> Notably, all previous studies used a dichotomized mRS as outcome measure for regression analyses, which has less statistical power than an ordinal analysis, and may thus lead to false-negative results.

In accordance with previous studies, prior use of IVT was not associated with a higher percentage of successful reperfusion,<sup>8,9,18</sup> nor was it associated with a higher percentage of first-pass effect, a relatively new measure of successful thrombectomy.<sup>12</sup> Conversely, procedure times were shorter in the IVT+EVT group. This implies that it may have been more difficult to gain intracranial access in the EVT alone group. It further implies that IVT does not facilitate the procedure by softening the thrombus. The latter could also have been influenced by differences in stroke cause between the 2 groups (ie, caused by a higher percentage of atrial fibrillation in the EVT-alone group) potentially leading to different clot characteristics.<sup>19</sup> However, if that were the case, the differences in successful reperfusion and first-pass effect would have been larger. Also, this association between clot characteristics and cause remains unclear,<sup>19</sup> and we did account for this imbalance in our multivariable analysis. Further, it is possible that the higher percentage of oral anticoagulation use in the EVT-only group influenced the occurrence of successful reperfusion, despite our attempts to adjust for this imbalance. However, there is currently no evidence that oral anticoagulation use and successful reperfusion are associated with one another.

We found that patients in the IVT+EVT group more frequently had reperfusion before start of EVT (eTICI  $\geq$ 2B on

**Table 4.** Primary, Secondary,\* and Safety Outcomes Among Patients Treated With IVT+EVT Versus EVT Alone

	IVT+EVT (n=1161)	EVT (n=324)	P Value	(c)OR/β (95% CI)	Adjusted (c)OR/β (95% CI) <sup>†</sup>
<b>Primary outcome</b>					
mRS at 90 d, median (IQR) <sup>‡</sup>	3 (2–6)	4 (2–6)	<0.01	1.80 (1.43–2.26)	1.47 (1.10–1.96) <sup>††</sup>
<b>Secondary outcomes</b>					
mRS 0–2 at 90 d, n (%)	431/1061 (41)	86/299 (29)	<0.01	1.65 (1.25–2.17)	1.32 (0.85–1.87) <sup>††</sup>
ΔNIHSS, median (IQR) <sup>§</sup>	−4 (−9 to 0)	−3 (−8 to 1)	0.02	−0.9 (−1.9 to 0.2)	−1.5 (−2.6 to −0.3) <sup>††</sup>
Door-intervention center-to-groin-puncture time for transferred patients <sup>  </sup>	47 (31–69)	47 (30–71)	0.72	−0.8 (−7.2 to 5.7)	0.5 (−7.3 to 8.2) <sup>††</sup>
Door-intervention center-to-groin-puncture time for nontransferred patients <sup>  </sup>	105 (79–130)	94 (73–125)	0.08	10.5 (2.2–18.8)	9.5 (0.5–18.5) <sup>††</sup>
Procedure time <sup>#</sup>	62 (39–87)	68 (45–95)	<0.01	−5.6 (−9.9 to −1.3)	−6.2 (−11.0 to −1.3) <sup>††</sup>
Onset-to-last-contrast-bolus time <sup>**</sup>	265 (215–324)	277 (221–355)	<0.01	−19.6 (−29.5 to −9.7)	−3.5 (−12.8 to 5.9) <sup>††</sup>
First-pass effect <sup>**  </sup>	147/842 (17)	41/259 (16)	0.543	1.20 (0.83–1.74)	1.22 (0.79–1.90) <sup>§§</sup>
Reperfusion (eTICI ≥2B) before start of EVT, n (%)	97/1161 (8.4)	9/324 (2.8)	<0.01	3.19 (1.59–6.39)	3.14 (1.47–6.73) <sup>§§</sup>
Successful reperfusion post-EVT (eTICI ≥2B), n (%)	672/1143 (59)	175/321 (54)	0.17	1.19 (0.92–1.52)	1.05 (0.77–1.43) <sup>††</sup>
Post-EVT eTICI ≥2C, n (%)	456/1143 (40)	120/321 (37)	0.42	1.11 (0.86–1.43)	0.98 (0.72–1.33) <sup>††</sup>
<b>Safety outcomes</b>					
Mortality at 90 d, n (%)	275/1161 (24)	122/324 (38)	<0.01	0.51 (0.40–0.67)	0.58 (0.40–0.82) <sup>††</sup>
Severe extracranial hemorrhage, n (%)	28/1161 (2.4)	5/324 (1.5)	0.35	1.58 (0.60–4.12)	1.96 (0.66–5.81) <sup>   </sup>
Symptomatic ICH, n (%)	69/1161 (5.9)	17/324 (5.3)	0.64	1.14 (0.66–1.97)	1.20 (0.64–2.25) <sup>   </sup>

(c)OR indicates common odds ratio; eTICI, extended Thrombolysis in Cerebral Ischemia scale; EVT, endovascular treatment; ICH, intracranial hemorrhage; IQR, interquartile range; IVT, intravenous thrombolysis; mRS, modified Rankin Scale score; NIHSS, National Institutes of Health Stroke Scale; ΔNIHSS, NIHSS at 24 to 48 hours minus baseline NIHSS.

\*All times are in minutes—median (IQR).

<sup>†</sup>All analyses were adjusted for: age (y), baseline NIHSS, history of atrial fibrillation, diabetes mellitus, hypertension, ischemic stroke, myocardial infarction, pre-stroke mRS, prior use of anticoagulant medication, onset-to-first noncontrast CT (NCCT) time.

Missing: <sup>‡</sup>125; <sup>§</sup>167; <sup>||</sup>35; <sup>#</sup>96; <sup>\*\*</sup>156; <sup>\*\*</sup>87.

<sup>††</sup>Additionally adjusted for: baseline mean arterial pressure (MAP), occlusion location, collateral score, transfer from a primary stroke center, center.

<sup>‡‡</sup>Additionally adjusted for: occlusion location, transfer from a primary stroke center, center.

<sup>§§</sup>Additionally adjusted for: occlusion location, transfer from a primary stroke center.

<sup>|||</sup>Additionally adjusted for: baseline MAP, prior use of antiplatelet agents, Alberta Stroke Program Early CT Score.


<sup>†††</sup>In patients with at least 1 attempt at thrombectomy with a device (n=1101/1267).

first digital subtraction angiography) than in the EVT-alone group. Reperfusion before start of EVT in the ESCAPE (Endovascular Treatment for Small Core and Anterior Circulation Proximal Occlusion with Emphasis on Minimizing CT to Recanalization Times) trial and MR CLEAN trial was found in 4.8% and 3.7% of patients, respectively, compared with 8% in our study.<sup>20,21</sup> Our supplementary analysis showed that patients with reperfusion before EVT had more distally located occlusions, supporting the hypothesis that IVT is most effective in more distally located thrombi.<sup>22</sup> However, these numbers are small and there may be underreporting of those patients who recovered before EVT because these patients were not included in the MR CLEAN Registry. Moreover, results of the recent EXTEND-IA TNK (Tenecteplase versus Alteplase before Endovascular Therapy for Ischemic Stroke) study suggest that tenecteplase may be a more effective fibrinolytic drug than alteplase in patients with a large vessel occlusion.<sup>23</sup> Whether tenecteplase can replace alteplase as

the preferred drug for IVT requires further study.<sup>24</sup> Conversely, the use of IVT before EVT may also pose a higher risk of emboli migrating to a previously uninvolved territory or thrombus migration to more distal arterial branches that cannot be reached with EVT, which is associated with a worse prognosis.<sup>25</sup> In our study, embolization to a new territory and thrombus migration were not documented. However, the postprocedural eTICI score of ≥2C was similar in both groups. This finding suggests that IVT contributes mainly to faster recanalization, rather than causing thrombus migration.

The median door-to-groin-puncture time was 11 minutes longer in nontransferred patients in the IVT+EVT group, which indicates that administration of IVT might contribute to a small delay in start of EVT. This difference was shorter than the 32-minute mean delay in the STAR (Solitaire FR Thrombectomy for Acute Revascularization) trial.<sup>26</sup> This suggests that, with the increased experience in intervention centers, delayed start of EVT because of IVT administration has become less of an issue.





**Figure 2.** Distribution of the modified Rankin Scale score at 90 days in IVT+EVT group vs EVT-alone group (%)\*. acOR indicates adjusted common odds ratio; EVT, endovascular treatment; IVT, intravenous thrombolysis; mRS, modified Rankin Scale. \*The mRS 0 to 5 group contains 125 missing cases, whereas the mRS 6 group is complete. Therefore, this figure over-represents mortality in both groups.

Our study showed a reduction in mortality in favor of IVT+EVT versus EVT alone, which is in contrast with previous studies.<sup>18,27,28</sup> Conversely, 1 previous study reported a lower mortality in the EVT-alone group.<sup>8</sup> Importantly, the IVT+EVT group of this study was matched with patients who were eligible for IVT but were treated with EVT alone at the physician's discretion. Because the current standard of care in The Netherlands is to always give IVT except when contraindicated, we could not perform a similar analysis.

The occurrence of sICH and severe extracranial hemorrhage did not differ between both groups, which is in line with most previous studies.<sup>7,9,16,18</sup> In the HERMES (Highly Effective Reperfusion evaluated in Multiple Endovascular Stroke Trials) meta-analysis, the risk of sICH was similar in the IVT+EVT group and IVT alone group.<sup>1</sup> This suggests that the occurrence of sICH can be mainly attributed to IVT rather than to EVT. Therefore,

our results might be a hallmark of residual (ie, unmeasured) confounding.

The stratified analysis revealed that the benefit of IVT before EVT was not present in patients with pre-existing atrial fibrillation. While this sensitivity analysis must be interpreted with caution, it could imply that IVT has a lower treatment effect in patients with a large vessel occlusion caused by embolism from the heart than in patients with stroke of other etiology. For instance, it may be that cardiac thrombi have a different composition or age than thrombi of noncardiac origin.<sup>29</sup> Another potential explanation could be the risk of sICH. The proportion of patients who developed a sICH after EVT alone (ie, without prior IVT) was almost 5% higher in patients without atrial fibrillation than in patients with atrial fibrillation (7.3% versus 2.8%, Table 8), which could partly explain the worse outcome of patients without atrial fibrillation who received EVT alone. On the

**Table 5.** No Recanalization Versus Recanalization Before Start of EVT in Patients With and Without IVT Per Occlusion Location\*

	IVT+EVT		EVT	
	No Recanalization Before Start of EVT (n=1015)	Recanalization Before Start of EVT (n=85)	No Recanalization Before Start of EVT (n=299)	Recanalization Before Start of EVT (n=8)
Occlusion location, n (%)				
ICA	66 (94)	4 (5.7)	12 (100)	0 (0)
ICA-T	236 (98)	5 (2.1)	71 (100)	0 (0)
Proximal M1	273 (94)	15 (5.2)	71 (95)	4 (5.3)
Distal M1	310 (89)	39 (11)	111 (100)	0 (0)
M2/M3	125 (85)	22 (15)	34 (92)	3 (8.1)
A1/A2	5 (100)	0 (0)	0 (0)	1 (100)

A1 indicates segment 1 of the anterior cerebral artery; A2, segment 2 of the anterior cerebral artery; EVT, endovascular treatment; ICA, internal carotid artery; ICA-T, terminal internal carotid artery; IVT, intravenous thrombolysis; M1, segment 1 of middle cerebral artery; M2, segment 2 of middle cerebral artery; M3, segment 3 of middle cerebral artery.

\*Missing data on occlusion location: 66. We excluded 12 patients in whom no occlusion was visible on computed tomography angiography after adjudication by the imaging core laboratory.

**Table 6.** Baseline Characteristics After Propensity Score Matching

	IVT+EVT (n=305)	EVT (n=305)	P Value
Age (y), median (IQR)	72 (59–82)	72 (63–80)	0.61
Men, n (%)	165/305 (54)	163/305 (53)	0.87
NIHSS, median (IQR)	16 (11–20)	17 (13–20)	0.13
Systolic BP, mean mm Hg (SD)	149 (25)	149 (26)	0.74
Diastolic BP, mean mm Hg (SD)	82 (16)	82 (17)	0.99
Medical history			
Atrial fibrillation, n (%)	102/305 (33)	126/305 (41)	0.045
Diabetes mellitus, n (%)	61/305 (20)	51/305 (17)	0.30
Hypertension, n (%)	157/305 (52)	167/305 (55)	0.42
Ischemic stroke, n (%)	62/305 (20)	74/305 (24)	0.24
Myocardial infarction, n (%)	53/305 (17)	62/305 (20)	0.35
Peripheral artery disease, n (%)	31/305 (10)	35/305 (11)	0.60
Pre-stroke mRS, n (%)			0.06
0	192/305 (63)	161/305 (53)	
1	41/305 (13)	54/305 (18)	
2	23/305 (7.5)	36/305 (12)	
≥3	49/305 (16)	54/305 (18)	
Medication			
Direct oral anticoagulants, n (%)	10/305 (3.3)	23/305 (7.5)	0.02
Vitamin K antagonists, n (%)	66/305 (22)	107/305 (35)	<0.01
Antiplatelets, n (%)	97/305 (32)	98/305 (32)	0.93
Imaging			
Occlusion location on CTA, n (%)			0.65
ICA	16/305 (5.3)	12/305 (3.9)	
ICA-T	69/305 (23)	77/305 (25)	
M1	174/305 (57)	180/305 (59)	
M2	41/305 (13)	33/305 (11)	
Other*	5/305 (1.6)	3/305 (1.0)	
ASPECTS, median (IQR)	8 (7–10)	9 (7–10)	0.66
Collateral score, n (%)			0.82
Grade 0	20/305 (6.6)	24/305 (7.9)	
Grade 1	101/305 (33)	94/305 (31)	
Grade 2	120/305 (39)	117/305 (38)	
Grade 3	64/305 (21)	70/305 (23)	

Baseline characteristics of set 1 of 5 imputed data sets are provided. ASPECTS indicates Alberta Stroke Program Early CT Score; BP, blood pressure; CTA, computed tomography angiography; EVT, endovascular treatment; ICA, internal carotid artery; ICA-T, terminal internal carotid artery; IQR, interquartile range; IVT, intravenous thrombolysis; M1, middle cerebral artery segment 1; M2, middle cerebral artery segment 2; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale.

\*Other: occlusions in segment 1 or 2 of the anterior cerebral artery (A1: n=2; A2: n=2) or no occlusion visible (n=4) on CTA after adjudication by the imaging core laboratory.

other hand, the difference in response to IVT in patients with atrial fibrillation could also be the result of residual confounding.

Finally, it is important to consider the implications of withholding IVT in patients eligible for both IVT and EVT, in

whom EVT may be delayed or not feasible. This is the case, for example, when the clot is not accessible because of technical reasons such as arterial tortuosity and extracranial carotid stenosis or occlusion.<sup>30</sup> In the MR CLEAN trial and Registry, this occurred in 5% and 6% of all patients, respectively, as

**Table 7.** Baseline Characteristics Among Patients Treated With IVT+EVT Versus EVT Alone, Stratified by Past Medical History of Atrial Fibrillation

	Patients With Atrial Fibrillation			Patients Without Atrial Fibrillation		
	IVT+EVT (n=186)	EVT (n=141)	P Value	IVT+EVT (n=958)	EVT (n=179)	P Value
Age (y), median (IQR)	77 (68–84)	77 (69–83)	0.75	69 (57–78)	68 (58–76)	0.54
Men, n (%)	87/186 (47)	70/141 (50)	0.61	525/958 (55)	100/179 (56)	0.79
NIHSS, median (IQR)*	16 (12–21)	17 (13–20)	0.36	15 (11–19)	16 (12–20)	0.04
Systolic BP, mean mm Hg (SD) <sup>†</sup>	148 (130–165)	150 (132–166)	0.54	150 (131–165)	147 (130–165)	0.21
Diastolic BP, mean mm Hg (SD) <sup>‡</sup>	84 (70–95)	80 (70–94)	0.70	80 (70–90)	80 (66–92)	0.77
Medical history						
Diabetes mellitus, n (%)	34/186 (18)	30/139 (22)	0.46	158/958 (16)	25/178 (14)	0.41
Hypertension, n (%)	114/184 (62)	98/140 (70)	0.13	438/947 (46)	79/177 (45)	0.69
Ischemic stroke, n (%)	36/186 (19)	39/141 (28)	0.08	126/957 (13)	43/178 (24)	<0.001
Myocardial infarction, n (%)	27/184 (15)	30/136 (22)	0.09	134/949 (14)	33/176 (19)	0.11
Peripheral artery disease, n (%)	17/181 (9.4)	16/139 (12)	0.54	78/948 (8.2)	20/176 (11)	0.18
Pre-stroke mRS, n (%)			0.27			<0.001
0	111/185 (60)	72/140 (51)		707/938 (75)	91/176 (52)	
1	27/185 (15)	23/140 (16)		103/938 (11)	32/176 (18)	
2	18/185 (9.7)	12/140 (8.6)		48/938 (5.1)	26/176 (15)	
≥3	29/185 (16)	33/140 (24)		80/938 (8.5)	27/176 (15)	
Medication						
Direct oral anticoagulants, n (%)	7/183 (3.8)	22/138 (16)	<0.001	3/944 (0.3)	5/177 (2.8)	<0.01
Vitamin K antagonists, n (%)	52/183 (28)	83/141 (59)	<0.001	18/952 (1.9)	35/179 (20)	<0.001
Antiplatelets, n (%)	60/183 (33)	33/139 (24)	0.08	327/949 (35)	66/177 (37)	0.47
Imaging						
Occlusion location on CTA, n (%)			0.02			0.74
ICA	5/175 (2.9)	4/137 (2.9)		65/913 (7.1)	8/165 (4.8)	
ICA-T	30/175 (17)	31/137 (23)		208/913 (23)	38/165 (23)	
M1	104/175 (59)	91/137 (66)		522/913 (57)	93/165 (56)	
M2	35/175 (20)	10/137 (7.3)		106/913 (12)	23/165 (14)	
Other <sup>  </sup>	1/175 (0.6)	1/137 (0.7)		12/913 (1.3)	3/165 (1.8)	
ASPECTS, median (IQR) <sup>§</sup>	9 (7–10)	9 (7–10)	0.37	9 (7–10)	9 (7–10)	0.51
Collateral score, n (%)			0.98			0.33
Grade 0	14/168 (8.3)	10/132 (7.6)		56/893 (6.3)	14/167 (8.4)	
Grade 1	56/168 (33)	42/132 (32)		304/893 (34)	51/167 (31)	
Grade 2	70/168 (42)	58/132 (44)		341/893 (38)	58/167 (35)	
Grade 3	28/168 (17)	22/132 (17)		192/893 (22)	44/167 (26)	

ASPECTS indicates Alberta Stroke Program Early CT Score; BP, blood pressure; CTA, computed tomography angiography; EVT, endovascular treatment; ICA, internal carotid artery; ICA-T, terminal internal carotid artery; IVT, intravenous thrombolysis; IQR, interquartile range; M1, middle cerebral artery segment 1; M2, middle cerebral artery segment 2; mRS, modified Rankin Scale; NIHSS, National Institutes of Health Stroke Scale.

Missing: \*29; <sup>†</sup>40; <sup>‡</sup>45; <sup>§</sup>64.

<sup>||</sup>Other: occlusions in segment 1 or 2 of the anterior cerebral artery, segment 3 of the middle cerebral artery, or no occlusion visible on CTA after adjudication by the imaging core laboratory.

opposed to the other randomized controlled trials that used more strict selection criteria and that reported percentages

between 2% and 3.6%.<sup>20,21,31,32</sup> In such cases, IVT remains the only available reperfusion therapy.

**Table 8.** Functional and Safety Outcomes Among Patients Treated With IVT+EVT Versus EVT Alone, Stratified by Past Medical History of Atrial Fibrillation

	Patients With Atrial Fibrillation				Patients Without Atrial Fibrillation			
	IVT+EVT (n=186)	EVT (n=141)	(c)OR (95% CI)	Adjusted (c)OR (95% CI)*	IVT+EVT (n=958)	EVT (n=179)	(c)OR (95% CI)	Adjusted (c)OR (95% CI)*
mRS at 90 d, median (IQR) <sup>†</sup>	4 (2–6)	5 (2–6)	1.33 (0.89–1.98)	1.06 (0.63–1.81) <sup>‡</sup>	3 (2–5)	4 (2–6)	1.74 (1.31–2.31)	1.72 (1.23–2.42) <sup>‡</sup>
mRS 0–2 at 90 d, n (%)	52/174 (30)	33/131 (25)	1.28 (0.78–2.12)	0.78 (0.37–1.65) <sup>‡</sup>	373/872 (43)	53/166 (32)	1.62 (1.15–2.27)	1.53 (0.97–2.42) <sup>‡</sup>
Mortality at 90 d, n (%)	65/186 (35)	57/141 (40)	0.80 (0.51–1.25)	0.87 (0.46–1.64) <sup>‡</sup>	204/958 (21)	64/179 (36)	0.50 (0.35–0.70)	0.44 (0.28–0.68) <sup>‡</sup>
Symptomatic ICH, n (%)	11/186 (5.9)	4/141 (2.8)	2.3 (0.72–7.39)	2.18 (0.60–7.91) <sup>§</sup>	56/958 (5.8)	13/179 (7.3)	0.81 (0.43–1.51)	0.94 (0.46–1.92) <sup>§</sup>
Successful reperfusion post-EVT (eTICI ≥2B), n (%)	95/186 (51)	76/139 (55)	0.85 (0.55–1.31)	0.73 (0.44–1.21)	568/940 (60)	97/178 (55)	1.28 (0.93–1.76)	1.15 (0.80–1.66)

(c)OR indicates common odds ratio; eTICI, extended Thrombolysis in Cerebral Ischemia scale; EVT, endovascular treatment; ICH, intracranial hemorrhage; IQR, interquartile range; IVT, intravenous thrombolysis; mRS, modified Rankin Scale score.

\*All analyses were adjusted for: age (y), baseline National Institutes of Health Stroke Scale (NIHSS), diabetes mellitus, hypertension, ischemic stroke, myocardial infarction, pre-stroke mRS, prior use of anticoagulant medication, onset-to-first noncontrast CT (NCCT) time.

<sup>†</sup>Missing: 122.

<sup>‡</sup>Additionally adjusted for: baseline mean arterial pressure (MAP), occlusion location, collateral score, transfer from a primary stroke center, center.

<sup>§</sup>Additionally adjusted for: baseline MAP, prior use of antiplatelet agents, Alberta Stroke Program Early CT Score.

Whether patients with acute ischemic stroke who are eligible for EVT should still receive intravenous alteplase is currently heavily debated among stroke physicians and researchers. In the present study, we explored many aspects of this debate. In addition to comparing clinical and safety outcomes in patients with IVT+EVT to those with EVT alone, we also included a complete assessment of workflow and procedural outcomes. Other strengths of our study include its comprehensive statistical approach, the large sample size, adherence to current protocols (eg, not awaiting the effect of IVT), and central adjudication of neuro-imaging, outcomes, and complications. Moreover, the data come from a nationwide registry and therefore reflect routine clinical practice. Similar to other studies that have tried to examine the added benefit of IVT before EVT, the main limitation of our study is the fact that it was nonrandomized. Patients in the EVT-alone group were selected based on contraindications for IVT. We also found that patients in the EVT-alone group more often had a pre-stroke disability, had a higher NIHSS at baseline, and were more often presented outside the 4.5-hour time window for IVT (reflected in the 16 minutes shorter onset-to-first noncontrast-computed-tomography time in patients with IVT+EVT). These factors are all associated with a worse prognosis.<sup>33,34</sup> The apparent benefit of prior IVT on functional outcome remained after adjustment for these factors and after propensity score matching based on these variables, yielding comparable point estimates of 1.47 and 1.42, respectively, but not after stratification for atrial fibrillation. Therefore, we still cannot exclude the possibility of residual confounding. Secondly, we had no data on whether the full dose of alteplase was administered in each patient who was treated with IVT. However, the Dutch stroke guideline does not advise on halting infusion of alteplase after groin puncture and/or recanalization. Since this guideline is well adhered to in The Netherlands, patients most likely received the full dose of alteplase.<sup>35</sup> Thirdly, patients who had a sICH before EVT and those who recovered before EVT are not represented in our study, since both types of patients would not have received EVT and were thus not recorded in the Registry.

Our data show that, in daily clinical practice, administering IVT before EVT has advantages and disadvantages. Because this is a nonrandomized study, we cannot firmly conclude whether IVT before EVT is beneficial to patients with acute ischemic stroke caused by an intracranial large vessel occlusion. Patients treated with both IVT and EVT had a better functional outcome and lower mortality at 90 days than those with contraindications for IVT who were treated with EVT alone, also after adjustment for potential confounders and after propensity score matching. Notably, in patients with atrial fibrillation, IVT may not be beneficial before EVT. Our results may indicate a true added benefit of IVT before EVT in select patient groups, but the interpretation is hampered by the possibility

of residual confounding or selection bias, which cannot be overcome by multivariable regression analysis or propensity score matching. MR CLEAN-NO IV (ISRCTN10888758), SWIFT-DIRECT (NCT03192332), DIRECT-SAFE (NCT03494920), and DIRECT-MT (NCT03469206), the 4 ongoing randomized clinical trials that directly compare both treatment strategies, will provide conclusive results on this topic. Meanwhile, IVT should not be withheld in patients outside these trials who are eligible for both IVT and EVT.

## Appendix

### MR CLEAN Registry Investigators

Diederik W. J. Dippel, MD, PhD (Erasmus MC University Medical Center Rotterdam, Executive Committee; Chair Writing Committee); Aad van der Lugt, MD, PhD (Erasmus MC University Medical Center Rotterdam, Executive Committee; Chair Imaging Assessment Committee; Chair Writing Committee); Charles B. L. M. Majoie, MD, PhD (Amsterdam UMC, University of Amsterdam, Executive Committee; Local Principal Investigator; Chair Imaging Assessment Committee; Chair Writing Committee); Yvo B. W. E. M. Roos MD, PhD (Amsterdam UMC, University of Amsterdam, Executive Committee; Chair Writing Committee); Robert J. van Oostenbrugge, MD, PhD (Maastricht University Medical Center, Executive Committee; Chair Writing Committee; Chair Adverse Event Committee); Wim H. van Zwam, MD, PhD (Maastricht University Medical Center, Executive Committee; Local Principal Investigator; Chair Imaging Assessment Committee; Chair Writing Committee); Jelis Boiten, MD, PhD (Haaglanden Medisch Centrum, the Hague, Executive Committee; Local Principal Investigator; Writing Committee); Jan Albert Vos, MD, PhD (Sint Antonius Hospital Nieuwegein, Executive Committee; Local Principal Investigator; Imaging Assessment Committee; Writing Committee); Ivo G. H. Jansen, MD, PhD (Amsterdam UMC, University of Amsterdam, Study Coordinator); Maxim J. H. L. Mulder, MD, PhD (Erasmus MC University Medical Center Rotterdam, Study Coordinator); Robert-Jan B. Goldhoorn, MD (Maastricht University Medical Center, Study Coordinator); Wouter J. Schonewille, MD, PhD (Sint Antonius Hospital Nieuwegein, Local Principal Investigator; Writing Committee); Jonathan M. Coutinho, MD, PhD (Amsterdam UMC, University of Amsterdam, Local Principal Investigator); Marieke J. H. Wermer, MD, PhD (Leiden University Medical Center, Local Principal Investigator); Marianne A. A. van Walderveen, MD, PhD (Leiden University Medical Center, Local Principal Investigator; Chair Imaging Assessment Committee); Julie Staals, MD, PhD (Maastricht University Medical Center, Local Principal Investigator); Jeannette Hofmeijer, MD, PhD (Rijnstate Hospital Arnhem, Local Principal Investigator; Writing Committee; Adverse Event Committee); Jasper M.

Martens, MD (Rijnstate Hospital Arnhem, Local Principal Investigator; Imaging Assessment Committee; Writing Committee); Geert J. Lycklama à Nijeholt, MD, PhD (Haaglanden Medisch Centrum, the Hague, Local Principal Investigator; Chair Imaging Assessment Committee; Writing Committee); Bob Roozenbeek, MD, PhD (Erasmus MC University Medical Center Rotterdam, Local Principal Investigator); Bart J. Emmer, MD, PhD (Amsterdam UMC, University of Amsterdam, Local Principal Investigator; Imaging Assessment Committee); Sebastiaan F. de Bruijn, MD, PhD (HAGA Hospital the Hague, Local Principal Investigator); Lukas C. van Dijk, MD (HAGA Hospital the Hague, Local Principal Investigator); H. Bart van der Worp, MD, PhD (University Medical Center Utrecht, Local Principal Investigator; Writing Committee); Rob H. Lo, MD (University Medical Center Utrecht, Local Principal Investigator; Writing Committee); Ewoud J. van Dijk, MD, PhD (Radboud University Medical Center Nijmegen, Local Principal Investigator); Hieronymus D. Boogaarts, MD, PhD (Radboud University Medical Center Nijmegen, Local Principal Investigator); Paul L. M. de Kort, MD, PhD (Elisabeth-TweeSteden Hospital Tilburg, Local Principal Investigator); Jo J. P. Peluso, MD, PhD (Elisabeth-TweeSteden Hospital Tilburg, Local Principal Investigator); Jan S. P. van den Berg, MD, PhD (Isala Klinieken Zwolle, Local Principal Investigator); Boudewijn A. A. M. van Hasselt, MD (Isala Klinieken Zwolle, Local Principal Investigator); Leo A. M. Aerden, MD, PhD (Reinier de Graaf Gasthuis Delft, Local Principal Investigator); René J. Dallinga, MD (Reinier de Graaf Gasthuis Delft, Local Principal Investigator); Maarten Uyttenboogaart, MD, PhD (University Medical Center Groningen, Local Principal Investigator); Omid Eshghi, MD (University Medical Center Groningen, Local Principal Investigator); Tobien H. C. M. L. Schreuder, MD (Atrium Medical Center Heerlen, Local Principal Investigator); Roel J. J. Heijboer, MD (Atrium Medical Center Heerlen, Local Principal Investigator); Koos Keizer, MD, PhD (Catharina Hospital Eindhoven, Local Principal Investigator); Lonneke S. F. Yo, MD (Catharina Hospital Eindhoven, Local Principal Investigator; Imaging Assessment Committee); Heleen M. den Hertog, MD, PhD (Isala Klinieken Zwolle, Local Principal Investigator); Emiel J. C. Sturm, MD (Medical Spectrum Twente Enschede, Local Principal Investigator); Marieke E. S. Sprengers, MD, PhD (Amsterdam UMC, University of Amsterdam, Imaging Assessment Committee); Sjoerd F. M. Jenniskens, MD, PhD (Radboud University Medical Center Nijmegen, Imaging Assessment Committee); René van den Berg, MD, PhD (Amsterdam UMC, University of Amsterdam, Imaging Assessment Committee); Albert J. Yoo, MD (Texas Stroke Institute United States of America, Imaging Assessment Committee); Ludo F. M. Beenen, MD (Amsterdam UMC, University of Amsterdam, Imaging Assessment Committee); Stefan D. Roosendaal, MD, PhD (Amsterdam UMC, University of Amsterdam, Imaging Assessment Committee); Bas F. W. van der



Kallen, MD, PhD (Haaglanden Medisch Centrum, the Hague, Imaging Assessment Committee); Ido R. van den Wijngaard, MD (Haaglanden Medisch Centrum, the Hague, Imaging Assessment Committee); Adriaan C. G. M. van Es, MD, PhD (Erasmus MC University Medical Center Rotterdam, Imaging Assessment Committee); Joseph C. J. Bot, MD, PhD (Amsterdam UMC, Vrije Universiteit Amsterdam, Imaging Assessment Committee); Pieter-Jan van Doormaal, MD (Erasmus MC University Medical Center Rotterdam, Imaging Assessment Committee); H. Zwenneke Flach, MD (Isala Klinieken Zwolle, Adverse Event Committee); Hester F. Lingsma, PhD (Erasmus MC University Medical Center Rotterdam, Trial Methodologist); Naziha el Ghannouti (Erasmus MC University Medical Center Rotterdam, Local Trial Coordinator); Martin Sterrenberg (Erasmus MC University Medical Center Rotterdam, Local Trial Coordinator); Corina Puppels (Sint Antonius Hospital Nieuwegein, Local Trial Coordinator); Wilma Pellikaan (Sint Antonius Hospital Nieuwegein, Local Trial Coordinator); Rita Sprengers (Amsterdam UMC, University of Amsterdam, Local Trial Coordinator); Marjan Elfrink (Rijnstate Hospital Arnhem, Local Trial Coordinator); Joke de Meris (Haaglanden Medisch Centrum, the Hague, Local Trial Coordinator); Tamara Vermeulen (Haaglanden Medisch Centrum, the Hague, Local Trial Coordinator); Annet Geerlings (Radboud University Medical Center Nijmegen, Local Trial Coordinator); Gina van Vemde (Isala Klinieken Zwolle, Local Trial Coordinator); Tiny Simons (Atrium Medical Center Heerlen, Local Trial Coordinator); Cathelijnn van Rijswijk (Elisabeth-TweeSteden Hospital Tilburg, Local Trial Coordinator); Gert Messchendorp (University Medical Center Groningen, Local Trial Coordinator); Hester Bongenaar (Catharina Hospital Eindhoven, Local Trial Coordinator); Karin Bodde (Reinier de Graaf Gasthuis Delft, Local Trial Coordinator); Sandra Kleijn (Medical Spectrum Twente Enschede, Local Trial Coordinator); Jasmijn Lodico (Medical Spectrum Twente Enschede, Local Trial Coordinator); Hanneke Droste (Medical Spectrum Twente Enschede, Local Trial Coordinator); M. Wollaert (Maastricht University Medical Center, Local Trial Coordinator); D. Jeurissen (Maastricht University Medical Center, Local Trial Coordinator); Ernas Bos (Leiden University Medical Center, Local Trial Coordinator); Yvonne Drabbe (HAGA Hospital the Hague, Local Trial Coordinator); Marjan Elfrink (Rijnstate Hospital Arnhem, Local Trial Coordinator); Berber Zweedijk (University Medical Center Utrecht, Local Trial Coordinator); Mostafa Khalilzada (HAGA Hospital the Hague, Local Trial Coordinator); Esme Venema (Erasmus MC University Medical Center Rotterdam, PhD Student); Vicky Chalos (Erasmus MC University Medical Center Rotterdam, PhD Student); Kars C. J. Compagne (Erasmus MC University Medical Center Rotterdam, PhD Student); Ralph R. Geuskens (Amsterdam UMC, University of Amsterdam, Medical Student); Tim van Straaten (Radboud University Medical Center Nijmegen, Medical Student); Saliha

Ergezen (Erasmus MC University Medical Center Rotterdam, Medical Student); Roger R. M. Harmsma (Erasmus MC University Medical Center Rotterdam, Medical Student); Anouk de Jong (Erasmus MC University Medical Center Rotterdam, PhD Student); Wouter Hinsenveld (Sint Antonius Hospital Nieuwegein, PhD Student); Olvert A. Berkhemer (Erasmus MC University Medical Center Rotterdam and Amsterdam UMC, University of Amsterdam and Maastricht University Medical Center, PhD); Anna M. M. Boers (Amsterdam UMC, University of Amsterdam, PhD); P. F. C. Groot (Amsterdam UMC, University of Amsterdam, Medical Student); Marieke A. Mens (Amsterdam UMC, University of Amsterdam, Medical Student); Katinka R. van Kranendonk (Amsterdam UMC, University of Amsterdam, PhD Student); Kilian M. Treurniet (Amsterdam UMC, University of Amsterdam, PhD Student); Manon Kappelhof (Amsterdam UMC, University of Amsterdam, PhD Student); Manon L. Tolhuijsen (Amsterdam UMC, University of Amsterdam, PhD Student); Heitor Alves (Amsterdam UMC, University of Amsterdam, PhD Student).

## Acknowledgments

We would like to thank the MR CLEAN Registry Investigators.

## Sources of Funding

The authors received no funding for this study. The MR CLEAN Registry was partly funded by Toegepast Wetenschappelijk Instituut voor Neuromodulatie (TWIN) Foundation, Erasmus MC University Medical Center, Maastricht University Medical Center, and Amsterdam UMC, University of Amsterdam.

## Disclosures

Dr LeCouffe, Dr Treurniet, and Dr Coutinho are research coordinators for the MR CLEAN-NO IV trial (ISRCTN80619088). Dr Roos and Dr Majoie are principal investigators of the MR CLEAN-NO IV trial. Dr Chalos, Dr Dippel, Dr van der Lugt, Dr Uyttenboogaart, Dr Lingsma, and Dr Roozenbeek are members of the CONTRAST (Collaboration for New Treatments of Acute Stroke) Consortium. Erasmus MC University Medical Center Rotterdam received compensation from Stryker<sup>®</sup> for consultations by Dr Dippel, Dr van der Lugt, and from Bracco Imaging<sup>®</sup> for consultations by Dr Dippel. Dr Dippel also reports research grants from Dutch Heart Foundation, Dutch Brain Foundation, and unrestricted grants from AngioCare BV, Medtronic/Covidien/EV3<sup>®</sup>, MEDAC GmbH/LAMEPRO, Penumbra Inc, Stryker<sup>®</sup>, and Top Medical/Concentric (all paid to the institution Erasmus MC University Medical Center Rotterdam). Dr van der Lugt also reports that Erasmus MC University Medical Center Rotterdam received unrestricted grants from

CVON/Dutch Heart Foundation, Dutch Brain Foundation, Stryker, Medtronic, and Penumbra for the conduct of studies for acute ischemic stroke and acute intracerebral hemorrhage. Amsterdam UMC, University of Amsterdam received compensation from Stryker® for consultations by Dr Majoie and Dr Roos. Dr Majoie also reports that Amsterdam UMC, University of Amsterdam received unrestricted grants from CVON/Dutch Heart Foundation, European Commission, TWIN Foundation, and Stryker. Dr Majoie and Dr Roos are shareholders of Nicolab. UMC Utrecht received grants from the Dutch Heart Foundation and compensation from Boehringer Ingelheim for consultations by Dr Van der Worp. The remaining authors have no disclosures to report.

## References

- Goyal M, Menon BK, van Zwam WH, Dippel DWJ, Mitchell PJ, Demchuk AM, Dávalos A, Majoie CBLM, van der Lugt A, de Miquel MA, Donnan GA, Roos YBWEM, Bonafe A, Jahan R, Diener H-C, van den Berg LA, Levy EI, Berkhemer OA, Pereira VM, Rempel J, Millan M, Davis SM, Roy D, Thornton J, Roman LS, Ribo M, Beumer D, Stouch B, Brown S, Campbell BC, van Oostenbrugge RJ, Saver JL, Hill MD, Jovin TG. Endovascular thrombectomy after large-vessel ischemic stroke: a meta-analysis of individual patient data from five randomised trials. *Lancet*. 2016;387:1723–1731.
- Powers WJ, Rabinstein AA, Ackerson T, Adeoye OM, Bambakidis NC, Becker K, Biller J, Brown M, Demaerschalk BM, Hoh B, Jauch EC, Kidwell CS, Leslie-Mazwi TM, Ovbiagele B, Scott PA, Sheth KN, Southerland AM, Summers DV, Tirschwell DL; American Heart Association Stroke Council on behalf of the AHA. 2018 guidelines for the early management of patients with acute ischemic stroke: a guideline for healthcare professionals from the American Heart Association/American Stroke Association. *Stroke*. 2018;49:e46–e110.
- Desilles J-P, Loyau S, Syvannarath V, Gonzalez-Valcarcel J, Cantier M, Louedec L, Lapergue B, Amarenco P, Ajzenberg N, Jandrot-Perrus M, Michel J-B, Ho-Tin-Noe B, Mazighi M. Alteplase reduces downstream microvascular thrombosis and improves the benefit of large artery recanalization in stroke. *Stroke*. 2015;46:3241–3248.
- Kidwell CS, Latour L, Saver JL, Alger JR, Starkman S, Duckwiler G, Jahan R, Vinuela F, Kang DW, Warach S. Thrombolytic toxicity: blood brain barrier disruption in human ischemic stroke. *Cerebrovasc Dis*. 2008;25:338–343.
- Fischer U, Kaesmacher J, Mendes Pereira V, Chapot R, Siddiqui AH, Froehler MT, Cognard C, Furlan AJ, Saver JL, Gralla J. Direct mechanical thrombectomy versus combined intravenous and mechanical thrombectomy in large-artery anterior circulation stroke. *Stroke*. 2017;48:2912–2918.
- Chandra RV, Leslie-Mazwi TM, Mehta BP, Derdeyn CP, Demchuk AM, Menon BK, Goyal M, González RG, Hirsch JA. Does the use of IV tPA in the current era of rapid and predictable recanalization by mechanical embolectomy represent good value? *J Neurointerv Surg*. 2016;8:443–446.
- Abilleira S, Ribera A, Cardona P, Rubiera M, López-Cancio E, Amaro S, Rodríguez-Campello A, Camps-Renom P, Cánovas D, de Miquel MA, Tomasello A, Remollo S, López-Rueda A, Vivas E, Perendreu J, Gallofré M; Catalan Stroke Code and Reperfusion Consortium. Outcomes after direct thrombectomy or combined intravenous and endovascular treatment are not different. *Stroke*. 2017;48:375–378.
- Broeg-Morvaj A, Mordasini P, Bernasconi C, Buhlmann M, Pult F, Arnold M, Schroth G, Jung S, Mattle HP, Gralla J, Fischer U. Direct mechanical intervention versus combined intravenous and mechanical intervention in large artery anterior circulation stroke: a matched-pairs analysis. *Stroke*. 2016;47:1037–1044.
- Coutinho JM, Liebeskind DS, Slater L-A, Nogueira RG, Clark W, Dávalos A, Bonafe A, Jahan R, Fischer U, Gralla J, Saver JL, Pereira VM. Combined intravenous thrombolysis and thrombectomy vs thrombectomy alone for acute ischemic stroke: a pooled analysis of the SWIFT and STAR studies. *JAMA Neurol*. 2017;74:268–274.
- Bellwald S, Ralph W, Arnold M, Von Martial R, Bernasconi C, Mordasini P, Chapot R, Fischer U, Gralla J. Direct mechanical intervention versus bridging therapy in large artery anterior circulation stroke: a pooled analysis of two registries. *Eur Stroke J*. 2017;2:302–303.
- Jansen IGH, Mulder MJHL, Goldhoorn RJB; MR CLEAN Registry Investigators. Endovascular treatment for acute ischaemic stroke in routine clinical practice: prospective, observational cohort study (MR CLEAN Registry). *BMJ*. 2018;360:k949.
- Zaidat OO, Castonguay AC, Linfante I, Gupta R, Martin CO, Holloway WE, Mueller-Kronast N, English JD, Dabus G, Malisch TW, Marden FA, Bozorgchami H, Xavier A, Rai AT, Froehler MT, Badruddin A, Nguyen TN, Taqi MA, Abraham MG, Yoo AJ, Janardhan V, Shaltoni H, Novakovic R, Abou-Chebl A, Chen PR, Britz GW, Sun C-HJ, Bansal V, Kaushal R, Nanda A, Nogueira RG. First pass effect: a new measure for stroke thrombectomy devices. *Stroke*. 2018;49:660–666.
- Goyal M, Fargen KM, Turk AS, Mocco J, Liebeskind DS, Frei D, Demchuk AM. 2C or not 2C: defining an improved revascularization grading scale and the need for standardization of angiography outcomes in stroke trials. *J Neurointerv Surg*. 2014;6:83–86.
- Von Kummer R, Broderick JP, Campbell BCV, Demchuk A, Goyal M, Hill MD, Treumiet KM, Majoie CBLM, Marquering HA, Mazya MV, Román LS, Saver JL, Strbian D, Whiteley W, Hacke W. The Heidelberg Bleeding Classification: classification of bleeding events after ischemic stroke and reperfusion therapy. *Stroke*. 2015;46:2981–2986.
- Tan IYL, Demchuk AM, Hopyan J, Zhang L, Gladstone D, Wong K, Martin M, Symons SP, Fox AJ, Aviv RI. CT angiography clot burden score and collateral score: correlation with clinical and radiologic outcomes in acute middle cerebral artery infarct. *AJNR Am J Neuroradiol*. 2009;30:525–531.
- Alonso de Leciana M, Martínez-Sánchez P, García-Pastor A, Kawiorski MM, Calleja P, Sanz-Cuesta BE, Díaz-Otero F, Frutos R, Sierra-Hidalgo F, Ruiz-Ares G, Fandiño E, Díez-Tejedor E, Gil-Núñez A, Fuentes B. Mechanical thrombectomy in patients with medical contraindications for intravenous thrombolysis: a prospective observational study. *J Neurointerv Surg*. 2017;9:1041–1046.
- Merlino G, Sponza M, Petralia B, Vit A, Gavrilovic V, Pellegrin A, Rana M, Cancelli I, Naliato S, Lorenzetti S, Marini R, Calzolari F, Eleopra R. Short and long-term outcomes after combined intravenous thrombolysis and mechanical thrombectomy versus direct mechanical thrombectomy: a prospective single-center study. *J Thromb Thrombolysis*. 2017;44:203–209.
- Weber R, Nordmeyer H, Hadisurya J, Heddier M, Stauder M, Stracke P, Berger K, Chapot R. Comparison of outcome and interventional complication rate in patients with acute stroke treated with mechanical thrombectomy with and without bridging thrombolysis. *J Neurointerv Surg*. 2017;9:229–233.
- Brinjikji W, Duffy S, Burrows A, Hacke W, Liebeskind D, Majoie CBLM, Dippel DWJ, Siddiqui AH, Khatri P, Baxter B, Nogueira R, Gounis M, Jovin T, Kallmes DF. Correlation of imaging and histopathology of thrombi in acute ischemic stroke with etiology and outcome: a systematic review. *J Neurointerv Surg*. 2017;9:529–534.
- Berkhemer OA, Fransen PSS, Beumer D, van den Berg LA, Lingsma HF, Yoo AJ, Schonewille WJ, Vos JA, Nederkoorn PJ, Wermer MJH, van Walderveen MAA, Staals J, Hofmeijer J, van Oostayen JA, Lycklama de Nijholt GJ, Boiten J, Brouwer PA, Emmer BJ, de Bruijn SF, van Dijk LC, Kappelle LJ, Lo RH, van Dijk EJ, de Vries J, de Kort PLM, van Rooij WJJ, van den Berg JSP, van Hasselt BAAM, Aerden LAM, Dallinga RJ, Visser MC, Bot JCI, Vroomen PC, Eshghi O, Schreuder THCM, Heijboer RJJ, Keizer K, Tielbeek AV, den Hertog HM, Gerrits DG, van den Berg-Vos RM, Karas GB, Steyerberg EW, Flach HZ, Marquering HA, Sprengers MES, Jenniskens SF, Beenen LFM, van den Berg R, Koudstaal PJ, van Zwam WH, Roos YBWEM, van der Lugt A, van Oostenbrugge RJ, Majoie CBLM, Dippel DWJ. A randomized trial of intraarterial treatment for acute ischemic stroke (MR CLEAN). *N Engl J Med*. 2015;372:11–20.
- Goyal M, Demchuk AM, Menon BK, Eesa M, Rempel JL, Thornton J, Roy D, Jovin TG, Willinsky RA, Sapkota BL, Dowlatshahi D, Frei DF, Kamal NR, Montaner WJ, Poppe AY, Ryckborst KJ, Silver FL, Shuaib A, Tampieri D, Williams D, Bang OY, Baxter BW, Burns PA, Choe H, Heo JH, Holmstedt CA, Jankowitz B, Kelly M, Linares G, Mandzia JL, Shankar J, Sohn SI, Swartz RH, Barber PA, Coutts SB, Smith EE, Morrish WF, Weill A, Subramanian S, Mitha AP, Wong JH, Lowerison MW, Sajobi TT, Hill MD. Randomized assessment of rapid endovascular treatment of ischemic stroke. *N Engl J Med*. 2015;372:1019–1030.
- Bhatia R, Hill MD, Shobha N, Menon B, Bal S, Kocher P, Watson T, Goyal M, Demchuk AM. Low rates of acute recanalization with intravenous recombinant tissue plasminogen activator in ischemic stroke: real-world experience and a call for action. *Stroke*. 2010;41:2254–2258.
- Campbell BCV, Mitchell PJ, Churilov L, Yassi N, Kleinig TJ, Dowling RJ, Yan B, Bush SJ, Dewey HM, Thijs V, Scroop R, Simpson M, Brooks M, Asadi H, Wu TY, Shah DG, Wijeratne T, Ang T, Miteff F, Levi CR, Rodrigues E, Zhao H, Salvaris P, Garcia-Esperon C, Bailey P, Rice H, de Villiers L, Brown H, Redmond K, Leggett D, Fink JN, Collicutt W, Wong AA, Muller C, Coulthard A, Mitchell K, Clouston J, Mahady K, Field D, Ma H, Phan TG, Chong W, Chandra RV, Slater L-A, Krause M, Harrington TJ, Faulder KC, Steinfort BS, Bladin CF, Sharma G, Desmond PM, Parsons MW, Donnan GA, Davis SM; EXTEND-IA TNK Investigators. Tenecteplase versus alteplase before thrombectomy for ischemic stroke. *N Engl J Med*. 2018;378:1573–1582.
- Hill MD, Michel P. Tenecteplase knocking on the door. *Stroke*. 2018;49:2276–2277.

25. Kaesmacher J, Maegerlein C, Kaesmacher M, Zimmer C, Poppert H, Friedrich B, Boeckh-Behrens T, Kleine JF. Thrombus migration in the middle cerebral artery: incidence, imaging signs, and impact on success of endovascular thrombectomy. *J Am Heart Assoc*. 2017;6:e005149. DOI: 10.1161/JAHA.116.005149.
26. Menon BK, Almekhlafi MA, Pereira VM, Gralla J, Bonafe A, Davalos A, Chapot R, Goyal M; STAR Study Investigators. Optimal workflow and process-based performance measures for endovascular therapy in acute ischemic stroke: analysis of the Solitaire FR thrombectomy for acute revascularization study. *Stroke*. 2014;45:2024–2029.
27. Kaesmacher J, Kleine JF. Bridging therapy with i. v. rtPA in MCA occlusion prior to endovascular thrombectomy: a double-edged sword? *Clin Neuroradiol*. 2018;28:81–89.
28. Kass-Hout T, Kass-Hout O, Mokin M, Thesier DM, Yashar P, Orion D, Jahshan S, Hopkins LN, Siddiqui AH, Snyder KV, Levy EI. Is bridging with intravenous thrombolysis of any benefit in endovascular therapy for acute ischemic stroke? *World Neurosurg*. 2014;82:e453–e458.
29. De Meyer SF, Andersson T, Baxter B, Bendszus M, Brouwer P, Brinjikji W, Campbell BC, Costalat V, Davalos A, Demchuk A, Dippel D, Fiehler J, Fischer U, Gilvarry M, Gounis MJ, Gralla J, Jansen O, Jovin T, Kallmes D, Khatri P, Lees KR, Lopez-Cancio E, Majoie C, Marquering H, Narata AP, Nogueira R, Ringleb P, Siddiqui A, Szikora I, Vale D, von Kummer R, Yoo AJ, Hacke W, Liebeskind DS. Analyses of thrombi in acute ischemic stroke: a consensus statement on current knowledge and future directions. *Int J Stroke*. 2017;12:606–614. DOI: 10.1177/1747493017709671.
30. Hong JT, Kim TH, Kim IS, Yang SH, Sung JH, Son BC, Lee SW. The effect of patient age on the internal carotid artery location around the atlas. *J Neurosurg Spine*. 2010;12:613–618.
31. Campbell BC, Mitchell PJ, Kleinig TJ, Dewey HM, Churilov L, Yassi N, Yan B, Dowling RJ, Parsons MW, Oxley TJ, Wu TY, Brooks M, Simpson MA, Miteff F, Levi CR, Krause M, Harrington TJ, Faulder KC, Steinfort BS, Priglinger M, Ang T, Scroop R, Barber PA, McGuinness B, Wijeratne T, Phan TG, Chong W, Chandra RV, Bladin CF, Badve M, Rice H, de Villiers L, Ma H, Desmond PM, Donnan GA, Davis SM. Endovascular therapy for ischemic stroke with perfusion-imaging selection. *N Engl J Med*. 2015;372:1009–1018.
32. Saver JL, Goyal M, Bonafe A, Diener H-C, Levy EI, Pereira VM, Albers GW, Cognard C, Cohen DJ, Hacke W, Jansen O, Jovin TG, Mattle HP, Nogueira RG, Siddiqui AH, Yavagal DR, Baxter BW, Devlin TG, Lopes DK, Reddy VK, du Mesnil de Rochemont R, Singer OC, Jahan R. Stent-retriever thrombectomy after intravenous t-PA vs. t-PA alone in stroke. *N Engl J Med*. 2015;372:2285–2295.
33. Linfante I, Starosciak AK, Walker GR, Dabus G, Castonguay AC, Gupta R, Sun C-HJ, Martin C, Holloway WE, Mueller-Kronast N, English JD, Malisch TW, Marden FA, Bozorgchami H, Xavier A, Rai AT, Froehler MT, Badruddin A, Nguyen TN, Taqi MA, Abraham MG, Janardhan V, Shaltoni H, Novakovic R, Yoo AJ, Abou-Chebl A, Chen PR, Britz GW, Kaushal R, Nanda A, Issa MA, Nogueira RG, Zaidat OO. Predictors of poor outcome despite recanalization: a multiple regression analysis of the NASA registry. *J Neurointerv Surg*. 2016;8:224–229.
34. Fransen PSS, Berkhemer OA, Lingsma HF, Beumer D, van den Berg LA, Yoo AJ, Schonewille WJ, Vos JA, Nederkoorn PJ, Wermer MJH, van Walderveer MAA, Staals J, Hofmeijer J, van Oostayen JA, Lycklama A Nijeholt GJ, Boiten J, Brouwer PA, Emmer BJ, de Bruijn SF, van Dijk LC, Kappelle LJ, Lo RH, van Dijk EJ, de Vries J, de Kort PLM, van den Berg JSP, van Hasselt BAAM, Aerden LAM, Dallinga RJ, Visser MC, Bot JCJ, Vroomen PC, Eshghi O, Schreuder THCM, Heijboer RJ, Keizer K, Tielbeek AV, den Hertog HM, Gerrits DG, van den Berg-Vos RM, Karas GB, Steyerberg EW, Flach HZ, Marquering HA, Sprengers MES, Jenniskens SFM, Beenen LFM, van den Berg R, Koudstaal PJ, van Zwam WH, Roos YBWEM, van Oostenbrugge RJ, Majoie CBLM, van der Lugt A, Dippel DWJ. Time to reperfusion and treatment effect for acute ischemic stroke: a randomized clinical trial. *JAMA Neurol*. 2016;73:190–196.
35. Dippel DWJ, van der Worp HB, Hofmeijer J, van den Berg-Vos RM, van Dijk EJ, Geurts M, Mess W, de Lau LML, Kouwenhoven M, Bouma BJ, de Borst GJ, Claassen JAH, van Eijk M, van den Born BJH, Boogaarts HD, van Zwam W, Brummer I, Kwakkel G, Verburg AFE, Kanselaar K, Deddens GJ, van Heugten CM, Schiemanck SK, Harbers A; Nederlandse Vereniging voor Neurologie. Richtlijn herseninfarct/-bloeding. The Netherlands. 2017:1–1715. Available at: [https://richtlijndatabase.nl/richtlijn/herseninfarct\\_en\\_hersenbloeding/startpagina\\_herseninfarct\\_bloeding.html](https://richtlijndatabase.nl/richtlijn/herseninfarct_en_hersenbloeding/startpagina_herseninfarct_bloeding.html). Accessed August 30, 2018.